

# Leaf hydraulic conductance, measured in situ, declines and recovers daily: leaf hydraulics, water potential and stomatal conductance in four temperate and three tropical tree species

D.M. JOHNSON,<sup>1,2</sup> D.R. WOODRUFF,<sup>1</sup> K.A. MCCULLOH<sup>3</sup> and F.C. MEINZER<sup>1</sup>

<sup>1</sup> USDA Forest Service, Pacific Northwest Research Station, Corvallis, OR 97331, USA

<sup>2</sup> Corresponding author (danieljohnson@fs.fed.us)

<sup>3</sup> Department of Wood Science and Engineering, Oregon State University, Corvallis, OR 97331, USA

Received February 8, 2009; accepted April 15, 2009; published online May 8, 2009

**Summary** Adequate leaf hydraulic conductance ( $K_{\text{leaf}}$ ) is critical for preventing transpiration-induced desiccation and subsequent stomatal closure that would restrict carbon gain. A few studies have reported midday depression of  $K_{\text{leaf}}$  (or petiole conductivity) and its subsequent recovery in situ, but the extent to which this phenomenon is universal is not known. The objectives of this study were to measure  $K_{\text{leaf}}$ , using a rehydration kinetics method, (1) in the laboratory (under controlled conditions) across a range of water potentials to construct vulnerability curves (VC) and (2) over the course of the day in the field along with leaf water potential and stomatal conductance ( $g_s$ ). Two broadleaf (one evergreen, *Arbutus menziesii* Pursh., and one deciduous, *Quercus garryana* Dougl.) and two coniferous species (*Pinus ponderosa* Dougl. and *Pseudotsuga menziesii* [Mirbel]) were chosen as representative of different plant types. In addition,  $K_{\text{leaf}}$  in the laboratory and leaf water potential in the field were measured for three tropical evergreen species (*Protium panamense* (Rose), *Tachigalia versicolor* Standley and L.O. Williams and *Vochysia ferruginea* Mart) to predict their daily changes in field  $K_{\text{leaf}}$  in situ. It was hypothesized that in the field, leaves would close their stomata at water potential thresholds at which  $K_{\text{leaf}}$  begins to decline sharply in laboratory-generated VC, thus preventing substantial losses of  $K_{\text{leaf}}$ . The temperate species showed a 15–66% decline in  $K_{\text{leaf}}$  by midday, before stomatal closure. Although there were substantial midday declines in  $K_{\text{leaf}}$ , recovery was nearly complete by late afternoon. Stomatal conductance began to decrease in *Pseudotsuga*, *Pinus* and *Quercus* once  $K_{\text{leaf}}$  began to decline; however, there was no detectable reduction in  $g_s$  in *Arbutus*. Predicted  $K_{\text{leaf}}$  in the tropical species, based on laboratory-generated VC, decreased by 74% of maximum  $K_{\text{leaf}}$  in *Tachigalia*, but only 22–32% in *Vochysia* and *Protium*. The results presented here, from the previous work of the authors and from other published studies, were consistent with two different strategies regarding

daily maintenance of  $K_{\text{leaf}}$ : (1) substantial loss and subsequent recovery or (2) a more conservative strategy of loss avoidance.

**Keywords:** cavitation, embolism, photosynthesis, transpiration, xylem.

## Introduction

The water pathway from the plant stem to the sites of evaporation in the leaf is critically important for maintaining leaf water balance and allowing stomata to stay open, resulting in carbon capture. However, water transport in the leaf is vulnerable to desiccation-induced changes, including conduit embolism and collapse (e.g., Bucci et al. 2003, Nardini et al. 2003, Cochard et al. 2004, Woodruff et al. 2007, Johnson et al. 2009). Recent research has suggested that these changes may be partially due to variation at the molecular scale, including aquaporin gene expression and protein conformation (Cochard et al. 2007, Kaldenhoff et al. 2008), that result in reductions in leaf hydraulic conductance ( $K_{\text{leaf}}$ ). There has been increasing interest in leaf hydraulics over the past 5 years, and it has been determined that  $K_{\text{leaf}}$  generally declines with increasing water stress, but that the extent of the decline and the water potentials corresponding to the decline, vary from species to species, even within a particular habitat (Salleo et al. 2001, Brodrigg and Holbrook 2003, Hao et al. 2008). Our understanding of the mechanisms responsible for the desiccation-induced decline in  $K_{\text{leaf}}$  is still far from complete, and is complicated by the interacting effects of light, temperature and water status on  $K_{\text{leaf}}$ . Voicu et al. (2008), Scoffoni et al. (2008) and Sellin et al. (2008) have all found increases in  $K_{\text{leaf}}$  with increasing light, and Sack et al. (2004) observed increases in  $K_{\text{leaf}}$  that were greater than would be expected due only to the reduced viscosity of water with increasing temperature.

Even less well understood is the phenomenon of  $K_{\text{leaf}}$  recovery while still under negative pressures (Clearwater and Goldstein 2005, Nardini et al. 2008).

The vast majority of studies on the effects of dehydration on leaf hydraulics have been conducted with excised leaves in the laboratory (e.g., Brodribb and Holbrook 2003, Lo Gullo et al. 2003, Nardini et al. 2003, Trifilò et al. 2003, Cochard et al. 2004) and only a few studies have measured leaf hydraulic conductance in situ (Brodribb and Holbrook 2003, 2004, Sack et al. 2003, Aranda et al. 2005, Sellin and Kupper 2007). Even fewer studies have focused on diurnal patterns of leaf hydraulics in the field. Brodribb and Holbrook (2004) found that in *Simarouba glauca* DC.  $K_{\text{leaf}}$  decreased by about half by midday compared to pre-dawn values and recovered completely by the end of the day. They also found a close correspondence between  $K_{\text{leaf}}$  vulnerability curves (VC) generated in the laboratory and those generated from measurements in the field. Similarly, Bucci et al. (2003) observed daily decreases and recovery in petiole conductivity in two tropical tree species and also found that at a given value of leaf water potential, laboratory generated VC yielded values of petiole conductivity similar to those measured in field-grown plants.

The objectives of this study were to determine the extent to which different species lose and regain  $K_{\text{leaf}}$  over the course of a typical summer day and to determine if contrasting species show similar relationships between daily time courses of  $K_{\text{leaf}}$  and stomatal conductance. The authors also wanted to assess the robustness of the rehydration kinetics method for quickly determining  $K_{\text{leaf}}$  in the field. Finally, leaf hydraulic VC and leaf water potentials measured in the field were used to predict the daily course of  $K_{\text{leaf}}$  for other species in which  $K_{\text{leaf}}$  had not been directly measured in the field.

## Materials and methods

### Field sites and species

Two field sites were used: one near Corvallis, OR and the other in the Republic of Panama. The first field site was chosen based on proximity to the laboratory, the species that were available at the site, and because a previous study with baseline physiological data for the species studied here was performed over the summers of 2002 and 2003 (Davis 2005). The site was located in McDonald-Dunn Research Forest (44°38' N and 123°18' W) and all plants were located in a young forest (clear cut about 15 years prior, Davis 2005) and were within 300 m of each other. The Panamanian study site was in an old-growth forest in the Parque Nacional San Lorenzo on the Caribbean side of the Isthmus of Panama (9°17' N and 79°58' W). All measurements were carried out between August and September 2008 at the Oregon site, and between December and February 2003 and 2004 at the Panamanian site.

To represent different plant functional groups, one evergreen broadleaf (*Arbutus menziesii* Pursh.) and one deciduous broadleaf (*Quercus garryana* Dougl.) species were selected at the Oregon field site. Two coniferous species were also selected from the same site; one from the *Pinus* clade (long, needle-like leaves, with no plicate mesophyll; *Pinus ponderosa* Dougl.) and one from the *Larix-Pseudotsuga* clade (shorter, flat needles with differentiated palisade and spongy mesophyll; *Pseudotsuga menziesii* [Mirbel] Franco var. *menziesii*) (Esau 1977, Gernandt et al. 2008). In addition, three evergreen species from the tropical site were selected: *Protium panamense* (Rose) I.M. Johnst., *Tachigalia versicolor* Standley and L.O. Williams, and *Vochysia ferruginea* Mart.

### Leaf hydraulic conductance and vulnerability

Leaf hydraulic conductance ( $\text{mmol m}^{-2} \text{s}^{-1} \text{MPa}^{-1}$ ) was determined using a timed rehydration method described in Brodribb and Holbrook (2003), which is based on an analogy between rehydrating a leaf and recharging a capacitor:

$$K_{\text{leaf}} = C \ln(\Psi_o/\Psi_f)/t,$$

where  $C$  is the capacitance,  $\Psi_o$  is the leaf water potential before partial rehydration,  $\Psi_f$  is the leaf water potential after partial rehydration and  $t$  is the duration of rehydration. Branches about 30–50 cm long were collected from trees early in the morning before significant transpirational water loss and were transported to the laboratory, recut under water and allowed to rehydrate for at least 4 h. Measurements of leaf water potential ( $\Psi_L$ ) were conducted over the next 3 days on excised leaves/fascicles for initial values ( $\Psi_o$ ), and for final values after a period of rehydration of  $t$  seconds ( $\Psi_f$ ). Distilled water was used for rehydration of  $K_{\text{leaf}}$  samples, and water temperature was maintained between 21 and 23 °C.

For measurement of  $K_{\text{leaf}}$  in the field, branches (about 10–20 cm in length) were collected from trees, and leaves were then excised for determination of  $\Psi_o$ , with no equilibration time ( $\Psi$  for leaves on the same shoot typically varied by  $> 0.1$  MPa). Leaf samples from the same branch were then rehydrated for a period of  $t$  seconds and  $\Psi_f$  was measured. Distilled water was used for rehydration of  $K_{\text{leaf}}$  samples and all measurements took place in the shade. These measurements (both field and laboratory) were performed on individual leaves of *Quercus*, *Arbutus*, *Tachigalia*, *Protium* and *Vochysia*, small shoots ( $\sim 3$  cm long) of *Pseudotsuga* and fascicles (three needles each) of *Pinus*.

Field measurements of  $K_{\text{leaf}}$  along with corresponding measurements of  $\Psi_L$  and stomatal conductance were performed over 4 days in August and September 2008 (*Quercus* and *Pseudotsuga* on August 28th and September 3rd and *Arbutus* and *Pinus* on September 9th and September 16th).

All measurements were made on four to eight leaves from six preselected individual leaves about every 90 min from 530 to 600 h (predawn) until 1600–1630 h PDT. All individual leaves were in open areas, and fully sunlit branches/leaves were chosen for measurement (with the exception of predawn measurements). Both laboratory and field measurements of  $K_{\text{leaf}}$  were normalized by dividing each  $K_{\text{leaf}}$  value by the corresponding maximum  $K_{\text{leaf}}$  for each species and condition (field or laboratory; maximum  $K_{\text{leaf}}$  = mean of 10 highest values) and reported as relative  $K_{\text{leaf}}$ . This procedure resulted in several ( $\sim 4$ – $5$ ) individual values of  $K_{\text{leaf}}$  being greater than one for each species and site (see Figure 1).

Values of  $C$  were estimated from pressure–volume curves (Scholander et al. 1965, Tyree and Hammel 1972) using the methods described by Brodribb and Holbrook (2003). Briefly, the  $\Psi_L$  corresponding to turgor loss was estimated as the inflection point of the graph of  $\Psi_L$  versus relative water content (RWC). The slope of the curve before and following turgor loss provided  $C$  in terms of RWC ( $C_{\text{RWC}}$ ) for pre- and post-turgor loss, respectively. Five to six leaves of each species were used to construct pressure–volume curves and to estimate  $C$ .

Pressure–volume curves were conducted on individual leaves of the broadleaf species, small shoots (about 3 cm) of *Pseudotsuga* and on fascicles of three needles of *P. ponderosa*. Branch samples of about 30–50 cm, from the same individual leaves that were used for rehydration and measurement of  $K_{\text{leaf}}$ , were excised early in the morning and recut under water in the laboratory. Branches were allowed to rehydrate for at least 4 h before pressure–volume analyses were performed. Pressure–volume curves were created by plotting the inverse of  $\Psi_L$  against RWC with alternate determinations of fresh mass and  $\Psi_L$  repeated during slow dehydration of the twig on the laboratory bench until

values of  $\Psi_L$  exceeded the measuring range of the pressure chamber ( $-4.0$  MPa). Leaf water potential was measured using a pressure chamber (PMS Instrument Company, Corvallis, OR). For normalizing  $C$  on a leaf area basis, leaf areas for the broadleaf species and *Pseudotsuga* were obtained with a scanner and ImageJ Version 1.27 image analysis software (Abramoff et al. 2004, National Institute of Mental Health, Bethesda, MD) and needle areas for *P. ponderosa* were determined by multiplying mean needle lengths and circumferences ( $n = 6$  needles per species).

#### Leaf water potentials and stomatal conductance

Stomatal conductance ( $g_s$ ) was measured with a steady-state porometer (LI-1600, Li-Cor, Lincoln, NE) and leaf temperatures were measured concomitantly with a fine-wire thermocouple (located in the LI-1600 chamber). One-sided leaf areas of foliage from the porometer measurements were obtained with a scanner and ImageJ. Leaf water potential was measured using a pressure chamber on individual leaves of *Quercus*, *Arbutus*, *Tachigalia*, *Protium* and *Vochysia*, small shoots ( $\sim 3$  cm) of *Pseudotsuga* and fascicles (three needles each) of *Pinus*. Measurements of  $g_s$  and  $\Psi_L$  were conducted on the same dates and over the same time intervals as  $K_{\text{leaf}}$  measurements and were performed on five leaves of each species (per time interval).

#### Sampling and statistics

For each species,  $K_{\text{leaf}}$  data were grouped (binned) over water potential ranges of about 0.3 MPa (i.e.,  $-0.61$  to  $-0.90$  MPa,  $-0.91$  to  $-1.20$  MPa, etc.) with the exception of the first bin for each species, which corresponded to 0 to  $-0.6$  MPa (Figures 1 and 2). Each bin contained between two and 14 measurements and a total of 46–56




Figure 1. Leaf hydraulic conductance ( $K_{\text{leaf}}$ ) measured in the laboratory (closed circles) and in the field (open circles) for (A) *P. menziesii*, (B) *Q. garryana*, (C) *P. ponderosa* and (D) *A. menziesii*. Dashed lines represent mean bulk leaf turgor loss point (TLP) and vertical and horizontal error bars represent standard error.

Table 1. Species measured, percent of  $K_{\max}$  at daily  $\Psi_{\min}$  and the method used to determine percent  $K_{\max}$  (either measured directly, calculated by measuring transpiration and the difference in  $\Psi_L$  in transpiring and non-transpiring leaves, or interpreted from VC/acoustic emission curves (AEC) and  $\Psi_{\min}$ ).

Species	Reference	% $K_{\max}$	$\Psi_{\min}$	% $K_{\max}$ method
<i>Prunus mahaleb</i> L.	Kikuta et al. (1997)	13	-2.25	Interp from $\Psi_{\min}$ & AEC
<i>Phillyrea angustifolia</i> L.	Kikuta et al. (1997)	72	-1.88	Interp from $\Psi_{\min}$ & AEC
<i>Ilex aquifolium</i> L.	Kikuta et al. (1997)	45	-1.46	Interp from $\Psi_{\min}$ & AEC
<i>Lauris nobilis</i> L.	Kikuta et al. (1997)	15	-2.65	Interp from $\Psi_{\min}$ & AEC
<i>Quercus ilex</i> L.	Kikuta et al. (1997)	64	-1.75	Interp from $\Psi_{\min}$ & AEC
<i>Acer campestre</i> L.	Kikuta et al. (1997)	50	-1.61	Interp from $\Psi_{\min}$ & AEC
<i>Viburnum tinus</i> L.	Kikuta et al. (1997)	82	-1.29	Interp from $\Psi_{\min}$ & AEC
<i>Ceratonia siliqua</i> L.	Kikuta et al. (1997)	90	-1.80	Interp from $\Psi_{\min}$ & AEC
<i>C. alliodora</i>	Meinzer et al. (2004)	68	-2.45	Calc by $E/\Delta\Psi_L$
<i>S. morotoni</i>	Meinzer et al. (2004)	64	-1.70	Calc by $E/\Delta\Psi_L$
<i>S. glauca</i>	Brodribb and Holbrook (2004)	30	-1.55	Measured in situ
<i>Byrsonima crassifolia</i> HBK.	Brodribb and Holbrook (2006)	100	-1.3	Interp from $\Psi_{\min}$ & VC
<i>Rehdera trinervis</i> Moldenke	Brodribb and Holbrook (2006)	76	-1.2	Interp from $\Psi_{\min}$ & VC
<i>Tectaria confluens</i> (F.v.Muell. ex Bak.) Pic.Serm.	Brodribb and Holbrook (2006)	59	-1.15	Interp from $\Psi_{\min}$ & VC
<i>Genipa americana</i> L.	Brodribb and Holbrook (2006)	65	-0.85	Interp from $\Psi_{\min}$ & VC
<i>C. brasiliense</i>	Bucci et al. (2003)	35	-1.35	Measured (in petioles)
<i>S. macrocarpa</i>	Bucci et al. (2003)	37	-1.30	Measured (in petioles)
<i>Cercis siliquastrum</i> L.	Nardini et al. (2003)	86	-1.85	Interp from $\Psi_{\min}$ & VC
<i>Hymenae stignocarpa</i> Mart. Ex Hayne	Hao et al. (2008)	30	-2.13	Interp from $\Psi_{\min}$ & VC
<i>Aegiphila lhotzkiana</i> Cham.	Hao et al. (2008)	34	-1.04	Interp from $\Psi_{\min}$ & VC
<i>Myrsine guianensis</i> (Aubl.) Kuntze	Hao et al. (2008)	24	-1.54	Interp from $\Psi_{\min}$ & VC
<i>Styrax ferrugineus</i> Nees & Mart.	Hao et al. (2008)	29	-1.41	Interp from $\Psi_{\min}$ & VC
<i>Tapirira guianensis</i> Aubl.	FCM, unpublished	9	-1.98	Interp from $\Psi_{\min}$ & VC
<i>V. ferruginea</i>	Current study	78	-1.42	Interp from $\Psi_{\min}$ & VC
<i>P. panamense</i>	Current study	68	-1.67	Interp from $\Psi_{\min}$ & VC
<i>T. versicolor</i>	Current study	27	-2.02	Interp from $\Psi_{\min}$ & VC
<i>P. ponderosa</i>	Current study	33	-1.86	Measured in situ
<i>P. menziesii</i>	Current study	31	-2.00	Measured in situ
<i>Q. garryana</i>	Current study	77	-3.64	Measured in situ
<i>A. menziesii</i>	Current study	82	-2.15	Measured in situ
<i>Alnus rubra</i> Bong.	DMJ, unpublished	70	-2.06	Measured in situ

leaves were used for each curve, depending on the species. However, curves were fitted through non-binned data, which reduced the correlation coefficient, but reflected the truer fit of curves through the data. To reflect daily trends in field measurements, data for  $K_{\text{leaf}}$ ,  $\Psi_L$  and  $g_s$  were grouped according to time across 60 min intervals (Figures 3, 4 and 5). Best fit nonlinear (sigmoid;  $y = y_0 + [a/(1 + e^{(x-x_0)/b})]$ ) curves were fit through laboratory  $K_{\text{leaf}}$  data,  $R^2$  values were calculated and Student's  $t$  tests were used to compare the mean of laboratory versus field  $K_{\text{leaf}}$ .

## Results

Overall,  $K_{\text{leaf}}$  declined with decreasing  $\Psi_L$  in all species (Figures 1 and 2). Losses in  $K_{\text{leaf}}$  with decreasing water potential were similar in laboratory and field measurements, although there was greater variation in  $K_{\text{leaf}}$  at a given  $\Psi_L$  in the field (Figure 1). Maximum  $K_{\text{leaf}}$  was greater in the field than in the laboratory for all four species in

which it was measured under both conditions (Table 2) with a mean increase of 47% from laboratory to field. *Pseudotsuga* displayed the greatest difference between laboratory and field maximum  $K_{\text{leaf}}$  (88%), while *Arbutus* had the smallest difference (3%).

In the field, measured  $K_{\text{leaf}}$  increased from predawn values as light and temperature increased (600–800 h) in all of the temperate species, with a 27, 16, 14 and 7.6% increase from predawn values in *Pinus*, *Pseudotsuga*, *Arbutus* and *Quercus*, respectively. Mean leaf temperature and photosynthetically active radiation (PAR) increased over the same time period by 1.2, 0.33, 0.35 and 2.7 °C and 57, 28, 119 and 36  $\mu\text{mol m}^{-2} \text{s}^{-1}$  for *Pinus*, *Pseudotsuga*, *Arbutus* and *Quercus*, respectively ( $P < 0.05$  for all comparisons of leaf temperature and PAR at 0600 versus 0800 h). *Pseudotsuga*, *Pinus* and *Quercus* all displayed decreased midday  $K_{\text{leaf}}$  (about 66, 59 and 30% reductions, respectively), relative to maximum values (Figure 3) which closely corresponded to minimum  $\Psi_L$  for the day. However, *Arbutus* showed little decrease in  $K_{\text{leaf}}$  (about 15% reduction), even though  $\Psi_L$  dropped to about -2.0 MPa




Figure 2. Leaf hydraulic conductance ( $K_{\text{leaf}}$ ) measured in the laboratory for (A) *T. versicolor*, (B) *V. ferruginea* and (C) *P. panamense*. Dashed vertical lines represent mean bulk leaf TLP, horizontal bars represent observed ranges of water potentials in the field and vertical and horizontal error bars represent standard error.

by midday. Although a large percentage of  $K_{\text{leaf}}$  was lost in both *Pseudotsuga* and *Pinus* by midday, there was substantial recovery by 1600 h. Midday relative  $K_{\text{leaf}}$  was 0.43 and 0.44 for *Pseudotsuga* and *Pinus*, respectively, but increased to 0.70 and 0.87, respectively, by 1600 h. *Quercus*  $K_{\text{leaf}}$  had completely recovered to morning values by late afternoon, although the midday  $K_{\text{leaf}}$  depression in *Quercus* (29%) was not as great as in *Pinus* or *Pseudotsuga*.

Mean minimum daily  $\Psi_L$  ranged from  $-1.86 (\pm 0.07)$  MPa for *Pinus* to  $-3.64 (\pm 0.11)$  MPa for *Quercus* and corresponded closely to earlier data collected in 2002 and 2003




Figure 3. Leaf hydraulic conductance ( $K_{\text{leaf}}$ , open circles) and water potential ( $\Psi_{\text{leaf}}$ , closed circles) measured in the field for (A) *P. menziesii*, (B) *P. ponderosa*, (C) *Q. garryana* and (D) *A. menziesii*. Minimum  $\Psi_{\text{leaf}}$  (from Davis 2005), measured in the same area in 2002 and 2003, is represented by the 'x' symbol. Curves are best-fit regressions fit through leaf water potential data using a cubic function ( $y = y_0 + ax + bx^2 + cx^3$ ;  $R^2$  values were 0.96, 0.85, 0.92 and 0.84 for panels A–D, respectively, and  $P$  values for all regressions were  $< 0.01$ ) and vertical error bars represent standard error.

by Davis (2005, see Figure 3). Large losses in  $K_{\text{leaf}}$  were predicted at midday from the laboratory-generated VC based on the minimum  $\Psi_L$  measured in the field for all




Figure 4. Measured field leaf water potential ( $\Psi_{\text{leaf}}$ , A) and predicted relative hydraulic conductance ( $K_{\text{leaf}}$ , B) for *T. versicolor*, *V. ferruginea* and *P. panamense*. Vertical error bars represent standard error.

species except *Arbutus* (Figures 1–3). In fact, in *Pinus*, *Pseudotsuga* and *Tachigalia*, mean minimum  $\Psi_L$  in the field corresponded to about the lowest point on the  $y$ -axis of the

VC (the minimum  $K_{\text{leaf}}$ ). Based on these water potential data (and leaf VC),  $K_{\text{leaf}}$  was predicted to decrease to about 78, 68 and 26% of maximum  $K_{\text{leaf}}$  by midday for *Vochysia*, *Protium* and *Tachigalia*, respectively (Figure 4). In addition,  $g_s$  in *Pseudotsuga*, *Pinus* and *Quercus* began to decline 2–4 h after  $K_{\text{leaf}}$  had already started decreasing (see Figures 3 and 5).

## Discussion

The phenomenon of decreasing  $K_{\text{leaf}}$  with increasing water stress is well documented (e.g., Nardini et al. 2001, Bucci et al. 2003, Brodribb and Holbrook 2006, Woodruff et al. 2007). There are multiple potential mechanisms for dehydration-induced declines in  $K_{\text{leaf}}$ , including xylem cavitation, xylem implosion/deformation (Cochard et al. 2004), changes to downstream element conductivity (e.g., mesophyll) via changes in turgor (Brodribb and Holbrook 2003) and even biochemical-scale phenomena affecting the water permeability of cell membranes (Heidecker et al. 2003, Luu and Maurel 2005). However, recent work, using the independent methods of acoustic emission (Kikuta et al. 1997, Johnson et al. 2009) and cryo-SEM (Woodruff et al. 2007, Johnson et al. 2009), has shown that leaf xylem embolism is likely a critical element in the depressions in  $K_{\text{leaf}}$  associated with dehydration.

Few studies have characterized diel cycles of decline and recovery of  $K_{\text{leaf}}$  in situ. Bucci et al. (2003) found diurnal cycles of loss and recovery of hydraulic conductivity in petioles of *Schefflera macrocarpa* (C. & S) Seem. and *Caryocar brasiliense* Camb. In addition, Brodribb and Holbrook (2004) also observed decline and recovery of  $K_{\text{leaf}}$  over the course of a day and were able to predict loss




Figure 5. Stomatal conductance ( $g_s$ ) measured in the field for (A) *P. menziesii*, (B) *P. ponderosa*, (C) *Q. garryana* and (D) *A. menziesii*. Vertical error bars represent standard error and dashed lines represent the time corresponding to daily minimum  $\Psi_{\text{leaf}}$ . Numbers in parentheses represent the actual minimum values of  $\Psi_{\text{leaf}}$ .

Table 2. TLP, pre- and post-TLP capacitance ( $C$ ) and maximum leaf hydraulic conductance ( $K_{\text{leaf}}$ ) in both the laboratory and field. TLP and  $C$ -values from pressure–volume analyses. Numbers in parentheses are standard errors (\* and \*\* indicate significantly different field and laboratory maximum  $K_{\text{leaf}}$  at  $P < 0.01$  and  $0.001$ , respectively).

Species	TLP (MPa)	Pre-TLP $C$		Post-TLP $C$		Maximum $K_{\text{leaf}}$	
		mol m <sup>-2</sup> MPa <sup>-1</sup>				Field	Laboratory
		mmol m <sup>-2</sup> s <sup>-1</sup> MPa <sup>-1</sup>					
<i>Q. garryana</i>	-3.63(0.24)	0.33(0.10)	1.74(0.15)	8.56(0.07)	6.00(0.07)**		
<i>A. menziesii</i>	-2.40(0.20)	0.70(0.04)	2.73(0.15)	13.72(0.21)	13.35(0.10)*		
<i>P. menziesii</i>	-2.30(0.21)	0.44(0.01)	0.98(0.02)	7.40(0.08)	3.94(0.18)**		
<i>P. ponderosa</i>	-2.23(0.14)	0.55(0.02)	1.11(0.12)	8.15(0.17)	5.28(0.15)**		
<i>T. versicolor</i>	-2.39(0.08)	0.56(0.09)	2.35(0.46)	n/a	25.56(0.74)		
<i>V. ferruginea</i>	-2.23(0.08)	0.60(0.11)	2.52(0.30)	n/a	28.28(0.64)		
<i>P. panamense</i>	-2.57(0.08)	1.62(0.29)	6.80(0.73)	n/a	13.45(0.56)		

and recovery of  $K_{\text{leaf}}$  based on field leaf water potentials and laboratory-generated VC. The data presented here also show close agreement between laboratory and field  $K_{\text{leaf}}$  VC. Using laboratory-generated  $K_{\text{leaf}}$  VC and field leaf water potential measurements, it was predicted that *Tachigalia*, *Protium* and *Vochysia* all experience daily declines and recovery of  $K_{\text{leaf}}$ , although the degree of recovery in *Tachigalia* was small by the afternoon. Consistent with this, Meinzer et al. (2004) reported diurnal variation in  $K_{\text{leaf}}$  in the tropical species *Schefflera morototoni* (Aubl) and *Cordia alliodora* (Ruiz & Pav.) based on in situ measurements of branch transpiration and differences in  $\Psi_L$  of covered and exposed leaves. It should be noted, however, that the actual predicted percentage decrease and recovery of  $K_{\text{leaf}}$  in this study, based on laboratory-generated data, may be somewhat biased, due to potential differences in laboratory and field  $K_{\text{leaf}}$  (see Figure 1, panel A).

In this study,  $K_{\text{leaf}}$  began to recover from midday to late afternoon while leaves were still experiencing water potentials of less than  $-1$  MPa. This phenomenon of restoration of conductance while the xylem is still under tension has been observed in other species as well. For example, Bucci et al. (2003) observed afternoon recovery of conductivity in petioles, and Nardini et al. (2008) found complete recovery of sunflower vein embolism at  $-0.33$  MPa. There is much debate in the current literature concerning the mechanism(s) of recovery of conductance in plant organs experiencing negative xylem pressure (see Clearwater and Goldstein 2005, for review). Several hypotheses suggest that there may be mobilization of solutes to embolized conduits or regions adjacent to embolized conduits to induce refilling, and thus restore conductance (Grace 1993, Canny 1995, 1998, Hacke and Sperry 2003, Pickard 2003); however, more recently-proposed mechanisms implicate aquaporins or abscisic acid (ABA) or both as playing large roles in embolism recovery (Lovisolo and Schubert 2006, Lovisolo et al. 2008). Aquaporin expression is upregulated in many plants with increasing water stress (see Tyerman et al. 2002, for review) and it is thought that ABA may trigger expression of aquaporins during drought stress (Liu

et al. 1994, Malz and Sauter 1999). It has been hypothesized that ABA may have dual roles in embolism recovery/repair: (1) by maintaining stomatal closure, thus preventing transpiration (Lovisolo et al. 2008) and (2) by triggering aquaporin expression in tissues adjacent to embolized conduits, which could induce radial water flow into the dysfunctional xylem (Kaldenhoff et al. 2008).

In the current study, there were large differences between maximum  $K_{\text{leaf}}$  measured in the field and in the laboratory, especially in *Pseudotsuga* and *Pinus* (88% and 54% increase in field  $K_{\text{leaf}}$  measurements as compared to those performed in the laboratory). Additionally, midmorning (800 h)  $K_{\text{leaf}}$  was about 16% greater than predawn  $K_{\text{leaf}}$  in *Pinus*, *Quercus* and *Arbutus* (and potentially *Pseudotsuga*, although  $K_{\text{leaf}}$  appeared to decrease in *Pseudotsuga* at 0700 h), even though leaf water potential was more negative than at predawn. Leaf temperatures increased over the same time period by  $\sim 1.1$  °C and PAR increased by  $\sim 60$   $\mu\text{mol m}^{-2} \text{s}^{-1}$ . This trend in an early morning increase in conductivity has also been observed in petioles of two savanna tree species (Bucci et al. 2003), leaves of several temperate tree species (Lo Gullo et al. 2005) and leaves of sunflower (Tsuda and Tyree 2000). Leaf hydraulic conductance has previously been shown to increase with both increasing temperature and sunlight (e.g., Tyree et al. 2005, Scoffoni et al. 2008, Sellin et al. 2008, Voicu et al. 2008), which likely explains the observed early morning trends in  $K_{\text{leaf}}$  observed here, and potentially the differences in maximum  $K_{\text{leaf}}$  observed in the laboratory (equilibrated in the dark) and in the field (no equilibration). In addition, although absolute values of  $K_{\text{leaf}}$  differed between the field and laboratory in this study, their relative values were generally similar at the same water potential.

Maximum stomatal conductance (and photosynthesis) appears to be tightly coupled to maximum  $K_{\text{leaf}}$  across plant life forms (Sack et al. 2003, Brodribb et al. 2005, 2007, Sack and Holbrook 2006). However, stomatal behavior, as it relates to declining  $K_{\text{leaf}}$ , varies between species. In other words, different plants have different ‘safety margins’ or differences between water potential at stomatal closure and water potential at some degree of loss of  $K_{\text{leaf}}$  (see Sack




Figure 6. Frequency distribution of compiled data from this and other studies (see text for references). Relative leaf hydraulic conductance (either as measured, calculated from transpiration values and water potential differences between transpiring and non-transpiring leaves, or as predicted based on published VC and minimum leaf water potentials) at daily minimum leaf water potentials ( $n = 31$ ). Data were placed into bins representing intervals of 10% relative  $K_{leaf}$  (e.g., 0.00–0.09, 0.10–0.19).

and Holbrook 2006). In the current study, the more conservative *Quercus* and *Arbutus* appeared to have much larger safety margins than did either of the conifers, both of which lost a substantial portion of  $K_{leaf}$  before stomatal closure. Other examples of plants with both wide and narrow safety margins can be found (e.g., Nardini and Salleo 2000, Cochard et al. 2002, Bucci et al. 2003, Brodribb and Holbrook 2004, Woodruff et al. 2007). However, a survey of the literature, along with data from the current study, suggests that plants tend to fall in one of two groups (i.e., are bimodally-distributed, Figure 6): plants that retain the majority of their  $K_{leaf}$  at their minimum daily  $\Psi_L$  and plants that have lost the majority of their  $K_{leaf}$  at their minimum daily  $\Psi_L$ . There was no apparent association between the mode of regulation of minimum  $K_{leaf}$  and the phylogeny, leaf phenology, functional group or habitat type of the 31 species represented in Figure 6.

Based on the data presented here, the rehydration kinetics method was a reliable method for making many estimates of in situ  $K_{leaf}$  fairly quick. Although no equilibration was used in this study, an earlier study (Scoffoni et al. 2008) suggests that long equilibration times are not necessary for accurate measurements using this method. In addition, little variation in  $\Psi_L$  (typically no more than 0.1–0.2 MPa) was observed between leaves on the same branch in this study. However, *A. rubra*, an understory species experiencing frequent sunflecks, showed higher variability of  $K_{leaf}$  and no consistent diurnal trends, and there were larger variations in  $\Psi_L$  between leaves on the same branch (DMJ, unpublished data). Therefore, this method may only be suitable for certain species or species growing in certain environments (e.g., open areas with little patchiness in sunlight).

## Acknowledgments

This research was supported in part by NSF Grant IOB-0544470. The authors would like to thank Logan Barnart, Amaranta Valenzuela, Jean-Christophe Domec, Paula Campanello and Geneveva Gatti for their helpful assistance with field data collection. The authors are also grateful to Dr. Barbara Lachenbruch and the AvH plant biology group for critical discussion of ideas.

## References

- Abramoff, M.D., P.J. Magelhaes and S.J. Ram. 2004. Image processing with ImageJ. *Biophotonics Int.* 11:36–42.
- Aranda, I., L. Gil and J.A. Pardos. 2005. Seasonal changes in apparent hydraulic conductance and their implications for water use of European beech (*Fagus sylvatica* L.) and sessile oak [*Quercus petraea* (Matt.) Liebl] in South Europe. *Plant Ecology* 179:155–167.
- Brodribb, T.J. and N.M. Holbrook. 2003. Stomatal closure during leaf dehydration, correlation with other leaf physiological traits. *Plant Physiol.* 132:2166–2173.
- Brodribb, T.J. and N.M. Holbrook. 2004. Diurnal depression of leaf hydraulic conductance in a tropical tree species. *Plant Cell Environ.* 27:820–827.
- Brodribb, T.J. and N.M. Holbrook. 2006. Declining hydraulic efficiency as transpiring leaves desiccate: two types of response. *Plant Cell Environ.* 29:2205–2215.
- Brodribb, T.J., N.M. Holbrook, M.A. Zwieniecki and B. Palma. 2005. Leaf hydraulic capacity in ferns, conifers and angiosperms: impacts on photosynthetic maxima. *New Phytol.* 165:839–846.
- Brodribb, T.J., T.S. Feild and G.J. Jordan. 2007. Leaf maximum photosynthetic rate and venation are linked by hydraulics. *Plant Physiol.* 144:1890–1898.
- Bucci, S.J., F.G. Scholz, G. Goldstein, F.C. Meinzer and L. Da S.L. Sternberg. 2003. Dynamic changes in hydraulic conductivity in petioles of two savanna tree species: factors and mechanisms contributing to the refilling of embolized vessels. *Plant Cell Environ.* 26:1633–1645.
- Canny, M.J. 1995. A new theory for the ascent of sap-cohesion supported by tissue pressure. *Ann. Bot.* 75:343–357.
- Canny, M.J. 1998. Applications of the compensating pressure theory of water transport. *Am. J. Bot.* 85:897–909.
- Clearwater, M.J. and G. Goldstein. 2005. Embolism repair and long distance water transport. *In* *Vascular Transport in Plants*. Eds. N.M. Holbrook and M.A. Zwieniecki. Elsevier Academic Press, London, UK.
- Cochard, H., L. Coll, X. Le Rous and T. Améglio. 2002. Unraveling the effects of plant hydraulics on stomatal closure during water stress in walnut. *Plant Physiol.* 128: 282–290.
- Cochard, H., F. Froux, S. Mayr and C. Coutand. 2004. Xylem wall collapse in water-stressed pine needles. *Plant Physiol.* 134:401–408.
- Cochard, H., J.-S. Venisse, T.S. Barigah, N. Brunel, S. Harbette, A. Guilliot, M.T. Tyree and S. Sakr. 2007. Putative role of aquaporins in variable hydraulic conductance of leaves in response to light. *Plant Physiol.* 143:122–133.
- Davis, K.J. 2005. Comparison of the water relations characteristics of woody plants in western Oregon. M.S. Thesis. Oregon State University.
- Esau, K. 1977. *Anatomy of seed plants*. Wiley, New York, 576 p.
- Gernandt, D.S., S. Magallón, G. López, O.Z. Flores, A. Willyard and A. Listonk. 2008. Use of simultaneous analyses

- to guide fossil-based calibrations of Pinaceae phylogeny. *Int. J. Plant Sci.* 169:1086–1099.
- Grace, J. 1993. Refilling of embolized xylem. *In* *Water Transport in Plants Under Climate Stress*. Eds. M. Borghetti, J. Grace and A. Raschi. Cambridge University Press, Cambridge, UK.
- Hacke, U.G. and J.S. Sperry. 2003. Limits to xylem refilling under negative pressure in *Lauris nobilis* and *Acer negundo*. *Plant Cell Environ.* 26:303–311.
- Hao, G., W.A. Hoffmann, F.G. Scholz, S.J. Bucci, F.C. Meinzer, A.C. Franco, K. Cao and G. Goldstein. 2008. Stem and leaf hydraulics of congeneric tree species from adjacent tropical savanna and forest ecosystems. *Oecologia* 155:405–415.
- Heidecker, M., L.H. Wegner, K.-A. Binder and U. Zimmermann. 2003. Turgor pressure changes trigger characteristic changes in the electrical conductance of the tonoplast and the plasmalemma of the marine alga *Valonia utricularis*. *Plant Cell Environ.* 26:1035–1051.
- Johnson, D.M., F.C. Meinzer, D.R. Woodruff and K.A. McCulloh. 2009. Leaf xylem embolism, detected acoustically and by cryo-SEM, corresponds to decreases in leaf hydraulic conductance in four evergreen species. *Plant Cell Environ.* (in press).
- Kaldenhoff, R., M. Ribas-Carbo, J. Flexas Sans, C. Lovisolo, M. Heckwolf and N. Uehlein. 2008. Aquaporins and plant water balance. *Plant Cell Environ.* 31:658–666.
- Kikuta, S.B., M.A. Lo Gullo, A. Nardini, H. Richter and S. Salleo. 1997. Ultrasound acoustic emissions from dehydrating leaves of deciduous and evergreen trees. *Plant Cell Environ.* 20:1381–1390.
- Liu, Q., M. Umeda and H. Uchimiya. 1994. Isolation and expression analysis of two rice genes encoding the major intrinsic protein. *Plant Mol. Biol.* 26:2003–2007.
- Lo Gullo, M.A., A. Nardini, P. Trifilò and S. Salleo. 2003. Changes in leaf hydraulics and stomatal conductance following drought stress and irrigation in *Ceratonia siliqua* (Carob tree). *Physiol. Plant.* 117:186–194.
- Lo Gullo, M.A., A. Nardini, P. Trifilò and S. Salleo. 2005. Diurnal and seasonal variations in leaf hydraulic conductance in evergreen and deciduous trees. *Tree Physiol.* 25:505–512.
- Lovisolo, C. and A. Schubert. 2006. Mercury hinders recovery of shoot hydraulic conductivity during grapevine rehydration: evidence from a whole-plant approach. *New Phytol.* 172:469–478.
- Lovisolo, C., I. Perrone, W. Hartung and A. Schubert. 2008. An abscisic acid-related reduced transpiration promotes gradual embolism repair when grapevines are rehydrated after drought. *New Phytol.* 180:642–651.
- Luu, D.T. and C. Maurel. 2005. Aquaporins in a challenging environment: molecular gears for adjusting plant water status. *Plant Cell Environ.* 28:85–96.
- Malz, S. and M. Sauter. 1999. Expression of two PIP genes in rapidly growing internodes of rice is not primarily controlled by meristem activity or cell expansion. *Plant Mol. Biol.* 40:985–995.
- Meinzer, F.C., S.A. James and G. Goldstein. 2004. Dynamics of transpiration, sap flow and use of stored water in tropical forest canopy trees. *Tree Physiol.* 24:901–909.
- Nardini, A. and S. Salleo. 2000. Limitation of stomatal conductance by hydraulic traits: sensing or preventing xylem cavitation? *Trees* 15:14–24.
- Nardini, A., M.T. Tyree and S. Salleo. 2001. Xylem cavitation in the leaf of *Prunus laurocerasus* and its impact on leaf hydraulics. *Plant Physiol.* 125:1700–1709.
- Nardini, A., S. Salleo and F. Raimondo. 2003. Changes in leaf hydraulic conductance correlate with leaf vein embolism in *Cercis siliquastrum* L. *Trees* 17:529–534.
- Nardini, A., M. Ramani, E. Gortan and S. Salleo. 2008. Vein recovery from embolism occurs under negative pressure in leaves of sunflower (*Helianthus annuus*). *Physiol. Plant.* 133:755–764.
- Pickard, W.F. 2003. The riddle of root pressure. I. Putting Maxwell's demon to rest. *Funct. Plant Biol.* 30:121–134.
- Sack, L. and N.M. Holbrook. 2006. Leaf hydraulics. *Annu. Rev. Plant Biol.* 57:361–381.
- Sack, L., P.D. Cowan, N. Jaikummar and N.M. Holbrook. 2003. The 'hydrology' of leaves: coordination of structure and function in temperate wood species. *Plant Cell Environ.* 26:1343–1356.
- Sack, L., C.M. Streeter and N.M. Holbrook. 2004. Hydraulic analysis of water flow through leaves of sugar maple and red oak. *Plant Physiol.* 134:1824–1833.
- Salleo, S., M.A. Lo Gullo, F. Raimondo and A. Nardini. 2001. Vulnerability to cavitation of leaf minor veins: any impact on leaf gas exchange? *Plant Cell Environ.* 24:851–859.
- Scholander, P.F., H.T. Hammel, E.D. Bradstreet and E.A. Hemmington. 1965. Sap pressure in vascular plants. *Science* 148:339–346.
- Scoffoni, C., A. Pou, K. Aasamaa and L. Sack. 2008. The rapid light response of leaf hydraulic conductance: new evidence from two experimental methods. *Plant Cell Environ.* 31:1803–1812.
- Sellin, A. and P. Kupper. 2007. Temperature, light and leaf hydraulic conductance of little-leaf linden (*Tilia cordata*) in a mixed forest canopy. *Tree Physiol.* 27:679–688.
- Sellin, A., E. Öunapuu and P. Kupper. 2008. Effects of light intensity and duration on leaf hydraulic conductance and distribution of resistance in shoots of silver birch (*Betula pendula*). *Physiol. Plant.* 134:412–420.
- Trifilò, P., A. Gascò, F. Raimondo, A. Nardini and S. Salleo. 2003. Kinetics of recovery of leaf hydraulic conductance and vein functionality from cavitation-induced embolism in sunflower. *J. Exp. Bot.* 54:2323–2330.
- Tsuda, M. and M.T. Tyree. 2000. Plant hydraulic conductance measured by the high pressure flow meter in crop plants. *J. Exp. Bot.* 51:823–828.
- Tyerman, S.D., C.M. Niemietz and H. Bramley. 2002. Plant aquaporins: multifunctional water and solute channels with expanding roles. *Plant Cell Environ.* 25:173–194.
- Tyree, M.T. and H.T. Hammel. 1972. The measurement of the turgor pressure and the water relations of plants by the pressure-bomb technique. *J. Exp. Bot.* 23:267–282.
- Tyree, M.T., A. Nardini, S. Salleo, L. Sack and B. El Omari. 2005. The dependence of leaf hydraulic conductance on irradiance during HPFM measurements: any role for stomatal response? *J. Exp. Bot.* 56:737–744.
- Voicu, M.C., J. Zwiazek and M.T. Tyree. 2008. Light response of hydraulic conductance in bur oak (*Quercus macrocarpa*) leaves. *Tree Physiol.* 28:1007–1015.
- Woodruff, D.R., K.A. McCulloh, J.R. Warren, F.C. Meinzer and B. Lachenbruch. 2007. Impacts of tree height on leaf hydraulic architecture and stomatal control in Douglas-fir. *Plant Cell Environ.* 30:559–569.